


Proposed ways in which abnormal behaviour may change brain pathways and structure



References

1. Kargel C, Massau C, Weiss S, et al. Diminished functional connectivity on the road to child sexual abuse in pedophilia. *The journal of sexual medicine* 2015;12(3):783-95. doi: 10.1111/jsm.12819 [published Online First: 2015/01/24]
2. Hadley JA, Nenert R, Kraguljac NV, et al. Ventral tegmental area/midbrain functional connectivity and response to antipsychotic medication in schizophrenia. *Neuropsychopharmacology* 2014;39(4):1020-30. doi: 10.1038/npp.2013.305 [published Online First: 2013/10/30]
3. Sheffield JM, Barch DM. Cognition and resting-state functional connectivity in schizophrenia. *Neurosci Biobehav Rev* 2016;61:108-20. doi: 10.1016/j.neubiorev.2015.12.007 [published Online First: 2015/12/25]
4. Mukherjee P, Whalley HC, McKirdy JW, et al. Altered amygdala connectivity within the social brain in schizophrenia. *Schizophr Bull* 2014;40(1):152-60. doi: 10.1093/schbul/sbt086 [published Online First: 2013/07/16]
5. Kneer J, Borchardt V, Kargel C, et al. Diminished fronto-limbic functional connectivity in child sexual offenders. *J Psychiatr Res* 2018 doi: 10.1016/j.jpsychires.2018.01.012 [published Online First: 2018/03/14]
6. Yu Q, Allen EA, Sui J, et al. Brain connectivity networks in schizophrenia underlying resting state functional magnetic resonance imaging. *Current topics in medicinal chemistry* 2012;12(21):2415-25. [published Online First: 2013/01/03]